


Appl. No. Unassigned; Docket No. NL02 0571 US  
Amdt. dated 01-DEC-2005  
Preliminary Amendment

**Amendments to the Claims**

1. (ORIGINAL) A phenylethanediol derivative, characterized in that the phenylethanediol derivative comprises at least one photo-convertible group suitable for adjusting the helical twisting power of the phenylethanediol derivative.
2. (ORIGINAL) The phenylethanediol derivative of claim 1 further having at least one polymerizable group.
3. (CURRENTLY AMENDED) The phenylethanediol derivative of ~~claim 1~~ claim 1 wherein the photo-convertible group is a photo-isomerizable group.
4. (ORIGINAL) The phenylethanediol derivative of claim 3 wherein the photo-isomerizable group is an olefinic group.
5. (CURRENTLY AMENDED) The phenylethanediol derivative of ~~any one of claims 1-4~~ claim 1 wherein the polymerizable group is a (meth)acrylate group.
6. (CURRENTLY AMENDED) The phenylethanediol derivatives of ~~any one of the preceding claims~~ claim 1 wherein the phenylethanediol has the formula

Appl. No. Unassigned; Docket No. NL02 0571 US  
 Amdt. dated 01-DEC-2005  
 Preliminary Amendment




wherein

- A stands for a bond or a p-phenylene group;  
 B and B' are independently  $(O)_p-C_6H_{2o}-O-CO-CR'=CH_2$ , o being 2-12, p being 0 or 1,  
 and R' being H or  $CH_3$ ;  
 P stands for a  $CH_2$  or a  $C=O$  group;  
 Q and Q' are independently selected from H, C1-C3 alkyl, C1-C3 alkoxy, halogen,  
 and CN;  
 n is an integer from 1 to 3; and  
 m is an integer from 0 to 2.

7. (CURRENTLY AMENDED) The phenylethanedione derivative of any one  
 of the preceding claims claim 1 wherein the phenylethanedione has the formula

Appl. No. Unassigned; Docket No. NL02 0571 US  
 Amdt. dated 01-DEC-2005  
 Preliminary Amendment



wherein

A stands for a bond or a p-phenylene group;

B is  $(O)_p-C_6H_{2o}-O-CO-CR'=CH_2$ , o being 2-12, p is 1, and R' being H or  $CH_3$ ;

P stands for a  $CH_2$  or a  $C=O$  group;

Q is selected from H, C1-C3 alkyl, C1-C3 alkoxy, halogen, and CN; and  
 m is an integer from 0 to 2.

8. (ORIGINAL) A method for the preparation of the phenylethanedio derivative of claim 1 by the steps of a) synthesizing a 2-hydroxy ether-protected phenylethanedio, b) followed by etherification or esterification of the 1-hydroxy group of the 2-hydroxy ether-protected phenylethanedio with an alcohol (or derivative thereof) or acid, respectively, optionally comprising polymerizable and/or photo-convertible groups, c) then cleaving the ether-protective group to obtain a phenylethanedio derivative with a free 2-hydroxy group, and optionally d) esterification of the free 2-hydroxy group with an acid which optionally comprises one or more polymerizable and/or photo-convertible groups.

9. (CURRENTLY AMENDED) A cholesteric composition comprising the phenylethanedio derivative of ~~any one of claims 1-7~~ claim 1.

10. (CURRENTLY AMENDED) An optical element, preferably an optical color filter, comprising the phenylethanedio derivative of any one of claims 1-7.

11. (CURRENTLY AMENDED) Use of the phenylethanedio derivative of ~~any one of claims 1-7~~ claim 1 in optical elements.